

COMMON HYPERCYCLIC VECTORS FOR MULTIPLES OF BACKWARD SHIFT

EVGENY ABAKUMOV & JULIA GORDON

ABSTRACT. We prove that ℓ^2 contains a dense set of vectors which are hypercyclic simultaneously for all multiples of the backward shift operator by constants of absolute value greater than 1.

0. INTRODUCTION

The focus of our study in this paper is the hypercyclic behavior (density of an orbit) for scalar multiples of the backward shift operator.

For a linear topological space X , a continuous linear operator $T: X \rightarrow X$ is called *hypercyclic* if there exists a vector $x \in X$ such that its orbit $\{T^n x, n \geq 0\}$ is dense in X (such a vector x is called a *hypercyclic vector* for the operator T).

The study of the phenomenon of hypercyclicity originates in the papers by Birkhoff, 1929 [1] and MacLane, 1952 [7] that show, respectively, that the operators of translation and differentiation, acting on the space of entire functions, are hypercyclic. The first example of a hypercyclic operator acting on a Banach space was given by Rolewicz in 1969 [8]. The example is the backward shift operator on ℓ^p , scaled by a constant greater than 1.

Later on, starting from the eighties, the subject of hypercyclicity has been widely explored. We refer the reader to the paper by Godefroy and Shapiro [5] and the recent survey by Grosse-Erdmann [6] for information and references.

We will concentrate on the space ℓ^2 of all complex valued square summable sequences. The backward shift operator B on this space is defined by

$$B(a_0, a_1, a_2, \dots) = (a_1, a_2, a_3, \dots).$$

Since the operator B is a contraction, B itself cannot be hypercyclic. The above mentioned result due to Rolewicz states that the operator zB has a hypercyclic vector whenever $|z| > 1$. The question that we answer in this paper is whether it is possible to find a *single* vector $f \in \ell^2$ such that its orbit is dense in ℓ^2 for *all* operators zB with $|z| > 1$. The answer is *yes*:

Theorem. *Let B be the backward shift acting on ℓ^2 . There exists a common hypercyclic vector for the operators zB , $|z| > 1$.*

¹Keywords: hypercyclic vector, backward shift

²2000 *Mathematics Subject Classification*: 47A16, 47B37

For a continuous linear operator $T : X \rightarrow X$ acting on a Banach space X , the notation $HC(T)$ stands for the set of all vectors in X that are hypercyclic for T . It turns out that the set $HC(T)$ is residual (i.e. dense G_δ) whenever it is nonempty, see, e.g., [5]. This fact together with Baire Category Theorem immediately implies that if $\{T_\lambda, \lambda \in \Lambda\}$ is a family of hypercyclic operators, then $\cap_{\lambda \in \Lambda} HC(T_\lambda)$ is residual (in particular, nonempty) whenever Λ is *countable*.

For *uncountable* families of operators the standard Baire categories arguments fail, and this is what makes our question interesting. At the moment we do not know of any other non-trivial examples of uncountable families of operators with common hypercyclic vectors. The question about the set $\cap_{|z|>1} HC(zB)$ was asked by Salas [9]. Our theorem states that this set is nonempty, and hence dense (it suffices to notice that if x is a hypercyclic vector for an operator T , then each element of its orbit $\{T^n x, n \geq 0\}$ is also hypercyclic for T). We do not know if the set of common hypercyclic vectors is large or small (say, in the sense of Baire categories).

Let us mention here that analogous problems for *cyclic* vectors (those whose orbit *spans* a dense subset in the space) have been treated for some natural families of operators. In particular, there exists a common cyclic vector for adjoint multiplications on some spaces of analytic functions (see [10], [3], [2]), and for the collection of linear partial differential operators of finite order with constant coefficients (see [4]).

We will use the following notations: $\mathbb{N} = \{1, 2, \dots\}$ – the set of natural numbers; $\mathbb{R}_+$ – the set of positive real numbers; $[r]$ – the integral part of a real number, $[r] = \max\{n \in \mathbb{Z} \mid n \leq r\}$; B – the backward shift operator on the space ℓ^2 , as defined above; by $\|\cdot\|$ we will always mean ℓ^2 -norm.

Acknowledgement. We are grateful to Alexander Borichev for reading the manuscript and for his helpful suggestions.

1. ALGORITHM

We start with a review of the main constituents of our construction. For the simplicity of exposition, we construct a common hypercyclic vector for the family of real multiples $\{\lambda B, \lambda > 1\}$ of the operator B . Then, in Section 3.3, we describe how to modify the construction so that it would give a common hypercyclic vector for all complex multiples zB , $|z| > 1$.

1.1. Preliminary remarks. The first obvious observation is that if f is any vector in ℓ^2 and $n \in \mathbb{N}$ is fixed, then all the vectors $(\lambda B)^n f$, $\lambda > 0$, have the same direction as the vector $B^n f$. In particular, if we want f to be a common hypercyclic vector for the family $(\lambda B)_{\lambda > 1}$, then the sequence $B^n f$ should “approximate all directions”. To better define this requirement, we introduce a countable family of cones generated by a set of diminishing in size balls in ℓ^2 centered at the elements of a dense set in ℓ^2 . One of the requirements on a common hypercyclic vector f is that $B^n f$ falls into every cone sufficiently often.

To specify what “sufficiently often” means, we have to look closely at the norms of the vectors $B^n f$, where f is a common hypercyclic vector. Let us fix one of the cones $\mathcal{C}$ from the previous paragraph, and let $\{n_k\}_{k \geq 1}$ be the set of all integers n such that $B^n f \in \mathcal{C}$. Then, naturally, for all $\lambda > 1$ we have $(\lambda B)^n f \in \mathcal{C}$. Let $\mathcal{B}$ be a ball inside $\mathcal{C}$. Then for every $\lambda > 1$ there exists $k \in \mathbb{N}$ such that $\lambda^{n_k} B^{n_k} f \in \mathcal{B}$. In particular, this implies that for any $\lambda > 1$ there should exist an integer k such that the ℓ^2 -norm of the vector $\lambda^{n_k} B^{n_k} f$ belongs to a certain fixed interval $]a - \delta, a + \delta[\subset \mathbb{R}$. This turns out to be a very restrictive condition on the sequence of the norms $\{\|B^{n_k} f\|\}_{k \geq 1}$, as illustrated by the following statement.

Lemma. *Let $\{n_k\}_{k \geq 1}$ be a fixed increasing sequence of positive integers, and let $a \in \mathbb{R}_+$. Then the following conditions on a sequence $\{r_k\}_{k \geq 1}$ of real positive numbers are equivalent:*

- 1) *For every $\lambda > 1$, $\epsilon > 0$, there exists $k \in \mathbb{N}$ such that $|\lambda^{n_k} r_k - a| < \epsilon$.*
- 2) *Let $x_k = -\frac{\ln r_k}{n_k}$. Then for any $\delta > 0$, the $\frac{\delta}{n_k}$ -neighbourhoods of the points $\frac{\ln a}{n_k} + x_k$ cover $\mathbb{R}_+$.*

In particular, the sequence $\{\lambda^{n_k} r_k\}_{k \geq 1}$ is dense in $\mathbb{R}_+$ for any $\lambda > 1$ if and only if the condition (2) holds for a dense set of a 's.

Here the sequence $\{r_k\}_{k \geq 1}$ plays the role of $\{\|B^{n_k} f\|\}_{k \geq 1}$.

Proof. The proof is straightforward. $\square$

Remark. The lemma shows in particular that it is not always possible to find such a sequence $\{r_k\}_{k \geq 1}$ for a given sequence $\{n_k\}_{k \geq 1}$. For example, (2) implies that the Lebesgue measure of the union of $\frac{\delta}{n_k}$ -neighbourhoods of certain points is infinite; therefore, if $\{r_k\}_{k \geq 1}$ exists, then the series $\sum_{k \geq 1} \frac{1}{n_k}$ necessarily diverges.

It follows from this remark that if f is a common hypercyclic vector, and $\{n_k\}_{k \geq 1}$ is the sequence of all numbers such that the iterates $B^{n_k} f$ belong to a given cone, then the series $\sum_{k \geq 1} \frac{1}{n_k}$ must diverge, for every fixed cone. For this condition to hold, the iterates of the hypercyclic vector are required to “visit” every cone of our infinite family very often.

1.2. Main idea. Here is the algorithm that we follow in order to construct the common hypercyclic vector:

- 1) Fix a dense countable set in ℓ^2 .
- 2) Construct a sequence of cones generated by balls centered at the vectors of this set with radii going to 0. The sequence of all these cones will be denoted by $\{\mathcal{C}_l\}_{l=1}^\infty$.
- 3) Fix a function $j: \{n \in \mathbb{N}, n \geq 20\} \rightarrow \mathbb{N}$ that takes every positive integral value sufficiently often, and possesses certain additional properties.
- 4) Construct a sequence $\{M_n\}_{n \geq 20}$ of natural numbers simultaneously with a sequence $\{r_n\}_{n \geq 20}$ of positive real numbers. The sequence $\{M_n\}_{n \geq 20}$ will be the sequence of numbers of the iterates of the prospective common hypercyclic vector, over which we have complete control (see the

next step). This sequence will fall into a disjoint union of infinitely many subsequences according to the values of the function j : the l -th subsequence will consist of the elements of $\{M_n\}$ with $j(n) = l$. If we set $\{n_k\}_{k \geq 1}$ to be any of these subsequences without a few initial terms, and $\{r_k\}_{k \geq 1}$ to be the corresponding subsequence of the sequence $\{r_n\}_{n \geq 20}$, then $\{n_k\}_{k \geq 1}$ and $\{r_k\}_{k \geq 1}$ satisfy the condition (2) of the Lemma and some additional requirements.

- 5) Construct a vector $f \in \ell^2$ such that $\|B^{M_k} f\| = r_k$ for all $k \geq 20$, and such that for any $l \geq 1$, $B^{M_k} f$ belongs to the cone $\mathcal{C}_l$ for all sufficiently large k verifying $j(k) = l$.

It is roughly clear from the previous subsection that such a vector will be a common hypercyclic vector for the operators λB , $\lambda > 1$. We give a rigorous proof of this statement in Section 3. Section 2 is devoted to Step (4) of the algorithm, that is, to the construction of the sequences $\{M_k\}$ and $\{r_k\}$.

1.3. Notation. Here we carry out the Steps (1), (2), and (3) of the algorithm.

1.3.1. We call a vector $v \in \ell^2$ *finite*, if all its coordinates, except for finitely many, equal 0.

Let $\{v_l\}_{l \geq 1}$ be a dense subset of ℓ^2 , such that for all $l \in \mathbb{N}$ the vector v_l is nonzero and finite, and put $\alpha_l = \ln \|v_l\|$. Clearly, it is possible to choose the sequence $\{v_l\}_{l \geq 1}$ in such a way that the real numbers α_l satisfy the following condition:

$$(1) \quad |\alpha_{l+1} - \alpha_l| \leq 1 \text{ for all } l \geq 1; \quad \alpha_1 = \alpha_2 = 0.$$

We fix a sequence of positive real numbers $\{\epsilon_l\}_{l \geq 1}$ such that $\epsilon_l < \|v_l\|$ and $\epsilon_l \rightarrow 0$ as $l \rightarrow \infty$, and define the cone $\mathcal{C}_l$ by $\mathcal{C}_l = \{v \in \ell^2 \mid \exists \mu > 0: \|\mu v - v_l\| < \epsilon_l\}$.

1.3.2. We define the function $j: \{n \in \mathbb{N}, n \geq 20\} \rightarrow \mathbb{N}$ in the following way: let $j(k) = k + 1 - [n \ln \ln n]$, where n is the unique integer satisfying the inequality $[n \ln \ln n] \leq k < [(n+1) \ln \ln(n+1)]$. (Notice that $j(20) = 1$.) This function takes every positive integral value infinitely many times: its domain is subdivided into intervals, on each of which $j(k)$ increases with unit step from the value 1 at the beginning of the interval. The constant 20 appears in the construction due to the inequality $\ln \ln 20 > 1$ which will be used below.

We will use the following property of the function $j(k)$ which is easy to check:

$$(2) \quad j(k) < [\ln \ln k] + 3 \text{ for } k \geq 20.$$

Now we are ready to begin the actual construction.

2. MAIN CONSTRUCTION

Here we carry out Step (4) of the algorithm declared in the previous section. Let $\{\alpha_l\}_{l \geq 1}$ be the sequence of real numbers from Section 1.3.1.

We construct two sequences: $\{M_k\}_{k \geq 20}$ of positive integers and $\{x_k\}_{k \geq 20}$ of real numbers, that possess the following properties:

- i) $\{M_k\}_{k \geq 20}$ is strictly increasing and $M_{k+1} - M_k \rightarrow +\infty$ as $k \rightarrow \infty$.
- ii) $\{M_k x_k\}_{k \geq 20}$ is strictly increasing and $M_{k+1} x_{k+1} - M_k x_k \rightarrow +\infty$ as $k \rightarrow \infty$.
- iii) For any integer $l \geq 1$, $s \in \mathbb{R}_+$, $\delta > 0$, and $K > 0$, there exists an integer $k > K$ such that $j(k) = l$ and $|s - (x_k + \frac{\alpha_l}{M_k})| < \frac{\delta}{M_k}$.

Returning to the notation of 1.2, we set $r_k = e^{-M_k x_k}$. Here the properties (i) and (ii) are necessary in order for a vector from Step (5) of 1.2 to exist; the property (iii) is what will translate into the condition (2) of the Lemma.

2.1. Definition of M_k, x_k . We will inductively construct the sequence $\{M_k\}_{k \geq 20}$ simultaneously with an auxilliary sequence of real numbers $\{y_k\}_{k \geq 20}$. Later we will set $x_k = y_k - \frac{\alpha_{j(k)}}{M_k}$.

Step 1. Set $M_{20} = 1, y_{20} = 1$.

Now we turn to the description of the step number q ($q > 1$). In every step several consecutive terms M_n and y_n are defined; denote the number of the last term defined in the first $(q-1)$ steps by N . We shall always assume that $j(N) = 1$, that is, a step can end only when the function j returns to the value 1.

Step q . Set

$$(3) \quad M_{N+1} = q^2(M_N + 1), \quad y_{N+1} = \frac{2}{q}.$$

Denote the difference $M_{N+1} - M_N$ by d . Now let N_1 be the minimal integer such that $N_1 > N$ and $j(N_1) = 1$. Then we set

$$(4) \quad M_{k+1} = M_k + d + q[\ln \ln k], \quad k = N+1, N+2, \dots, N_1-1;$$

$$(5) \quad y_{k+1} = y_k, \quad k = N+1, N+2, \dots, N_1-2;$$

$$(6) \quad y_{N_1} = y_{N_1-1} + \frac{1}{qM_{N_1}}.$$

After that we repeat the procedure: find N_2 – the minimal integer greater than N_1 such that $j(N_2) = 1$, and define M_k by the formula (4) for $k = N_1, N_1+1, \dots, N_2-1$, while the y 's are kept equal to y_{N_1} , until at $k = N_2$ the sequence $\{y_k\}$ again “makes a step” of the length $\frac{1}{qM_{N_2}}$. In this fashion, we proceed to construct the terms with the indices up to N_3, N_4 , and so on until y_{N_m} becomes greater than q for some m . Then the Step number q ends, so M_{N_m} and y_{N_m} are the last terms constructed on the Step q . Notice that by construction $j(N_m) = 1$ as required. Below we will also need the observation that in any case $y_{N_m} < 2q$, since every time y increases,

it increases by a number less than 1, and we stop as soon as y_{N_m} becomes greater than q .

This would have completed the construction, but we need to show that the above procedure eventually terminates, that is, that y_k eventually does become greater than q . Here is a proof:

2.2. Proof of termination of the algorithm. The proof of existence of an integer N_m such that $y_{N_m} > q$ is carried out by contradiction. Suppose that $y_{N_i} \leq q$ for $i = 1, 2, \dots$. By definition of the sequence y_k , this means that the series

$$(7) \quad \sum_{\substack{k > N, \\ j(k)=1}} \frac{1}{qM_k}$$

converges. By the construction of the sequence M_k , we have $M_k \leq M_{N+1} + d(k - N - 1) + q(k - N - 1) \ln \ln(k - 1)$ for $k > N$. This implies that $M_k \leq Ck \ln \ln k$ for some constant C that depends on q and N but does not depend on k .

By definition of the function $j(n)$, the convergence of the series (7) is equivalent to the convergence of the series $\sum_{s \geq 20} \frac{1}{M_{[s \ln \ln s]}}$ (recall that at the moment we are assuming that the Step number q never terminates, that is, an infinite number of M_k 's are defined by the formula (4), so this series is supposedly well defined). Combining this statement with the estimate on M_k , we see that our assumption ultimately implies that the series

$$(8) \quad \sum_{s \geq 20} \frac{1}{s \cdot \ln \ln s \cdot \ln \ln(s \ln \ln s)}$$

converges, which is not the case, and we arrive at a contradiction. $\square$

The construction of the sequences $\{M_k\}_{k \geq 20}$ and $\{y_k\}_{k \geq 20}$ is now completed. Set

$$x_k = y_k - \frac{\alpha_{j(k)}}{M_k}, \quad k \geq 20.$$

2.3. Verification of the properties. We now turn to the verification of the required properties of these sequences.

The property (i) is immediate from the construction.

2.3.1. Property (ii). We need an estimate from below for the difference $M_{k+1}x_{k+1} - M_kx_k$. In view of the way in which the sequences $\{M_k\}_{k \geq 20}$, $\{x_k\}_{k \geq 20}$ were constructed, there are two cases.

Case 1. The index k is such that both M_k and M_{k+1} (and, accordingly, x_k and x_{k+1}) are defined within the same step.

We will treat this case keeping the notations introduced in the description of Step number q , $q > 1$. Since $y_{k+1} \geq y_k$ within Step q , we have

$$M_{k+1}x_{k+1} - M_kx_k \geq y_k(d + q[\ln \ln k]) - \alpha_{j(k+1)} + \alpha_{j(k)}.$$

Recall that $j(k) < [\ln \ln k] + 3$ by (2) and $|\alpha_{l+1} - \alpha_l| \leq 1$ by (1). These estimates together yield

$$\alpha_{j(k)} - \alpha_{j(k+1)} = \sum_{i=j(k)}^{j(k+1)-1} \alpha_i - \alpha_{i+1} \geq -(j(k+1) - j(k)) > -[\ln \ln(k+1)] - 2.$$

Combining all of the above with the inequality $d \geq q^2$ which follows from (3), we get

$$M_{k+1}x_{k+1} - M_kx_k > \frac{2}{q}(q^2 + q[\ln \ln k]) - [\ln \ln(k+1)] - 2 > 2q - 3.$$

Case 2. The index k happened to be a “boundary” index, that is, M_k is defined on the Step $q-1$, whereas M_{k+1} is the first element of the q -th step.

Assume that $q > 2$ (for $q = 2$, we have $M_{21}x_{21} - M_{20}x_{20} > 0$). Again, we keep the notation of the previous section. Say, $k = N$ – the last index of the Step $q-1$. Then, by definition (3), $M_{k+1} = q^2(M_k + 1)$, and $x_{k+1} = \frac{2}{q} - \frac{\alpha_{j(k+1)}}{M_{k+1}}$. Recall that $j(N) = 1$, $j(N+1) \leq 2$ by definition of the function j , and $\alpha_1 = \alpha_2 = 0$, see (1). Besides, we know that $q-1 < y_N < 2(q-1)$ since y_N was constructed on the Step $q-1$. Finally,

$$\begin{aligned} M_{k+1}x_{k+1} - M_kx_k &> \frac{2}{q}M_{k+1} - 2qM_k \\ &\geq 2 \left(\frac{1}{q}M_{k+1} - qM_k \right) = 2 \left(\frac{q^2M_k + q^2}{q} - qM_k \right) = 2q. \end{aligned}$$

We see that $M_{k+1}x_{k+1} - M_kx_k$ is always positive and increases to ∞ . $\square$

2.3.2. Property (iii). Let us fix the numbers $s \in \mathbb{R}_+$, $\delta > 0$, and $l, K \in \mathbb{N}$. Pick a number $Q > K$ such that $\frac{1}{Q} < \delta$ and $\frac{2}{Q} < s < Q$. Let $\{n_m\}_{m \geq 1}$ be the set of all positive integers n , such that $j(n) = l$. We will think of $\{n_m\}$ as an increasing sequence of positive integers. Observe that all the numbers M_{n_m} with sufficiently large m were defined on the steps with numbers $q > Q$. Let us consider all the points $y_{n_m} = x_{n_m} + \frac{\alpha_l}{M_{n_m}}$ which were constructed on the q -th step. Their $\frac{1}{qM_{n_m}}$ -neighbourhoods cover the interval $]\frac{2}{q}, q[$ by construction, and since the $\frac{\delta}{M_{n_m}}$ -neighbourhoods are even larger, s falls in one of them. $\square$

3. COMMON HYPERCYCLIC VECTOR

3.1. Construction of the vector. As announced in the previous section, we set $r_k = e^{-M_kx_k}$. By the property (ii), the sequence r_k is decreasing and it tends to 0 as $k \rightarrow \infty$.

We turn to the construction of a vector $f \in \ell^2$ such that $\|B^{M_k}f\| = r_k$ and $B^{M_k}f$ belongs to the cone $C_{j(k)}$ for all sufficiently large $k \in \mathbb{N}$.

Let $\{v_l\}_{l \geq 1}$ be the dense subset of ℓ^2 from 1.3.1. For an integer $k \geq 20$, let $d_k = \sqrt{r_k^2 - r_{k+1}^2}$, and let $w_k \in \ell^2$ be the vector whose first $M_{k+1} - M_k$

coordinates coincide with those of $v_{j(k)}$ and the coordinates starting from $M_{k+1} - M_k + 1$ equal 0. Then set

$$(9) \quad f = \sum_{k \geq 20} \frac{d_k}{\|w_k\|} S^{M_k} w_k,$$

where S is the forward shift operator, $S(a_0, a_1, a_2 \dots) = (0, a_0, a_1, \dots)$.

It is clear that $\|B^{M_k} f\| = (\sum_{i=k}^{\infty} d_i^2)^{1/2} = r_k$.

We now turn to show that for any $l \geq 1$, the vector $B^{M_k} f$ falls into the cone $\mathcal{C}_l$ for all sufficiently large k such that $j(k) = l$. First, observe that $\|B^{M_k} f - \frac{d_k}{\|w_k\|} w_k\| = r_{k+1}$ by the definition of f . Second, the property (ii) of M_k, x_k implies that $\frac{d_k^2}{r_{k+1}^2} = \frac{r_k^2 - r_{k+1}^2}{r_{k+1}^2} \rightarrow +\infty$. Combining these two facts, we get

$$(10) \quad \left\| \frac{B^{M_k} f}{d_k} - \frac{w_k}{\|w_k\|} \right\| = \frac{r_{k+1}}{d_k} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

The vectors v_k were assumed to be finite. Therefore, for a fixed l , for all sufficiently large k such that $j(k) = l$, the vectors v_l and w_k coincide, since $M_{k+1} - M_k \rightarrow \infty$ as $k \rightarrow \infty$. Thus (10) yields $\|\frac{\|v_l\|}{d_k} B^{M_k} f - v_l\| < \epsilon_l$, and hence $B^{M_k} f$ belongs to the cone $\mathcal{C}_{j(k)}$ for all sufficiently large k such that $j(k) = l$.

3.2. Proof of hypercyclicity for $\lambda > 1$. To show that f is a common hypercyclic vector for all operators λB with $\lambda > 1$, it suffices to verify that for any fixed integer $l \geq 1$ there exists $k \geq 20$ such that $j(k) = l$ and

$$(11) \quad \|(\lambda B)^{M_k} f - v_l\| < 3\epsilon_l.$$

Fix $l \geq 1$, and let K be a positive integer such that for all $n > K$ the vector $B^{M_n} f$ falls into the cone $\mathcal{C}_l$ whenever $j(n) = l$. The existence of K was proved in the previous subsection. Now the statement immediately follows from Lemma. Indeed, the property (iii) implies that the condition (2) of the Lemma is satisfied if we set $a = e^{\alpha_l} = \|v_l\|$ and let $\{n_k\}_{k \geq 1}$ be the sequence of all numbers M_n with $n > K$ and $j(n) = l$. This is precisely one of the subsequences discussed in Step 4 of the algorithm described in 1.2. Then by the Lemma for any $\lambda > 1$ there exists $k > K$ with $j(k) = l$ such that $|\lambda^{M_k} \|B^{M_k} f\| - \|v_l\|| < \epsilon_l$ (recall that $r_k = \|B^{M_k} f\|$). Since the vector $(\lambda B)^{M_k} f$ belongs to the cone generated by the ball of radius ϵ_l centered at v_l , and its norm differs from the norm of v_l by less than ϵ_l , the condition (11) is fulfilled. $\square$

3.3. Complex case. Let us now consider the complex multiples zB , $|z| > 1$, of the operator B . We write z in the trigonometric form $z = \lambda e^{2\pi i \theta}$ with $\lambda > 1$ and $\theta \in [0, 1[$.

We preserve the same notation as before, except for the function $j(k)$ which should be defined here by means of the sequence $n \ln \ln \ln n$ (and

not $n \ln \ln n$ as in Section 1.3.2). We fix a cone $\mathcal{C}_l$, exactly as before. For real λ 's it was necessary to establish that $\frac{\delta}{M_k}$ -neighbourhoods of the points $\frac{\ln \|v_l\| - \ln \|B^{M_k} f\|}{M_k}$ (with M_k such that $B^{M_k} f$ belongs to $\mathcal{C}_l$) cover $\mathbb{R}_+$ for any $\delta > 0$. We achieved this by constructing a “very dense” sequence $\{x_k\}$ in $\mathbb{R}_+$ (see Section 2.1) and then producing a vector f such that $x_k = -\ln \|B^{M_k} f\|$.

Now, in the same fashion as was previously done for the sequence $\{x_k\}$, we construct the sequence $\{(x_k, \theta_k)\}$ of elements of the set $\mathbb{R}_+ \times [0, 1[$. More precisely, let y_k be as in Section 2.1; we will be constructing (y_k, θ_k) simultaneously with M_k . Let us look back at “Step q”. We keep defining y_k 's and M_k 's by the same formulas. The only modification is that now before y_k is allowed to make a step forward (see the formula (6)), the θ -component will have to increase from 0 to 1 with the decreasing steps $\frac{1}{qM_{N_i}}$, also staying constant at its every value with only M 's changing until the function $j(n)$ completes a cycle and returns to the value 1. The same argument as in Section 2.2 shows that y_k is allowed to move forward each time once θ has completed the cycle. The proof that each step terminates essentially remains the same.

Given the sequences $\{M_k\}$ and $\{(x_k, \theta_k)\}$, we define the vector f by

$$(12) \quad f = \sum_k \frac{d_k}{\|w_k\|} e^{2\pi i \theta_k} (S^{M_k} w_k)$$

in the notation of 3.1. The proof of its hypercyclicity works in the same way as in the previous section, with the use of the fact that now the square $\frac{\delta}{M_k}$ -neighbourhoods of the points (y_k, θ_k) cover the half strip $\mathbb{R}_+ \times [0, 1[$.

In conclusion, we remark that the theorem remains true (with essentially the same proof) for the backward shift acting on the spaces ℓ^p , $1 \leq p < \infty$, and on the space c_0 of sequences that tend to 0.

REFERENCES

- [1] G. D. Birkhoff, *Démonstration d'un théorème élémentaire sur les fonctions entières*, C.R.Acad.Sci. Paris **189**(1929), 473-475.
- [2] P. Bourdon and J. H. Shapiro, *Spectral synthesis and common cyclic vectors*, Michigan Math. J. **37**(1990), 71-90.
- [3] K. C. Chan, *Common cyclic vectors for operator algebras on spaces of analytic functions*, Indiana Univ. Math. J. **37**(1988), 919-928.
- [4] K. C. Chan, *Common cyclic entire functions for partial differential operators*, Integral Equations Operator Theory **13**(1990), 132-137.
- [5] G. Godefroy and J. H. Shapiro, *Operators with dense, invariant, cyclic vector manifolds*, J. Funct. Anal. **98**(1991), 229-269.
- [6] K.-G. Grosse-Erdmann, *Universal families and hypercyclic operators*, Bull. Amer.Math. Soc., **36**(3) (1999), 345-381.
- [7] G. R. MacLane, *Sequences of derivatives and normal families*, J.Analyse Math. **2**(1952), 72-87.
- [8] S. Rolewicz, *On orbits of elements*, Studia Math., **32**(1969), 17-22.
- [9] H. N. Salas, *Supercyclicity and weighted shifts*, Studia Math., **135**(1) (1999), 55-73.

- [10] W. R. Wogen, *On some operators with cyclic vectors*, Indiana Univ. Math. J. **27**(1978), 163-171.

EQUIPE D'ANALYSE ET DE MATHÉMATIQUES APPLIQUÉES, UNIVERSITÉ DE MARNE-LA-VALLÉE, 5 BOULEVARD DESCARTES, CHAMPS-SUR-MARNE, 77454 MARNE-LA-VALLÉE CEDEX 2, FRANCE

E-mail address: `abakumov@math.univ-mlv.fr`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR MI 48109-1109, USA

E-mail address: `julygord@umich.edu`